

RESEARCH

Open Access

Random C^* -ternary algebras and applicationYeol Je Cho^{1,2}, Reza Saadati³ and Young-Oh Yang^{4*}

*Correspondence:

yangyo@cheju.ac.kr

⁴Department of Mathematics, Jeju National University, Jeju, 690-756, Korea

Full list of author information is available at the end of the article

Abstract

In this paper, we introduce the concept of random C^* -ternary algebras and consider some properties of them. As an application we approximate a random C^* -ternary algebra homomorphism in these spaces.

MSC: Primary 39B52; 17A40; secondary 46B03

Keywords: random C^* -ternary algebra; C^* -ternary algebra homomorphism; random complex Banach spaces

1 Introduction

Ternary algebraic operations were considered in the 19th century by several mathematicians, such as Cayley [1], who introduced the notion of cubic matrix which, in turn, was generalized by Kapranov *et al.* [2]. The simplest example of such a non-trivial ternary operation is given by the following composition rule:

$$\{a, b, c\}_{ijk} = \sum_{1 \leq l, m, n \leq N} a_{nil} b_{ljm} c_{mkn}$$

for each $i, j, k = 1, 2, \dots, N$.

Ternary structures and their generalization, the so-called n -ary structures, raise certain hopes in view of their applications in physics. Some significant applications are as follows (see [3, 4]):

- (1) The algebra of nonions generated by two matrices

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & \omega \\ \omega^2 & 0 & 0 \end{pmatrix},$$

where $\omega = e^{\frac{2\pi i}{3}}$, was introduced by Sylvester as a ternary analog of Hamilton's quaternions (see [5]).

- (2) The quark model inspired a particular brand of ternary algebraic systems. The so-called Nambu mechanics is based on such structures (see [6]).

2 Random C^* -ternary algebra

In the section, we adopt the usual terminology, notations and conventions of the theory of random C^* -ternary algebra.

Throughout this paper, Δ^+ is the space of distribution functions, that is, the space of all mappings $F: \mathbf{R} \cup \{-\infty, \infty\} \rightarrow [0, 1]$ such that F is left-continuous and non-decreasing on \mathbf{R} , $F(0) = 0$, and $F(+\infty) = 1$. D^+ is a subset of Δ^+ consisting of all functions $F \in \Delta^+$ for which $l^-F(+\infty) = 1$, where $l^-f(x)$ denotes the left limit of the function f at the point x , that is, $l^-f(x) = \lim_{t \rightarrow x^-} f(t)$. The space Δ^+ is partially ordered by the usual point-wise ordering of functions, i.e., $F \leq G$ if and only if $F(t) \leq G(t)$ for all t in \mathbf{R} . For example an element for Δ^+ is the distribution function ε_a given by $\varepsilon_a(t) = 0$, if $t \leq a$ and 1 if $t > a$.

The maximal element for Δ^+ in this order is the distribution function ε_0 (see [7–9]).

Definition 2.1 ([8]) A mapping $T: [0, 1] \times [0, 1] \rightarrow [0, 1]$ is called a *continuous triangular norm* (briefly, a continuous t -norm) if T satisfies the following conditions:

- (a) T is commutative and associative;
- (b) T is continuous;
- (c) $T(a, 1) = a$ for all $a \in [0, 1]$;
- (d) $T(a, b) \leq T(c, d)$ whenever $a \leq c$ and $b \leq d$ for all $a, b, c, d \in [0, 1]$.

Typical examples of continuous t -norms are $T_P(a, b) = ab$, $T_M(a, b) = \min(a, b)$, and $T_L(a, b) = \max(a + b - 1, 0)$ (the Lukasiewicz t -norm).

Definition 2.2 ([9]) A *random normed space* (briefly, RN-space) is a triple (X, μ, T) , where X is a vector space, T is a continuous t -norm and μ is a mapping from X into D^+ such that the following conditions hold:

- (RN1) $\mu_x(t) = \varepsilon_0(t)$ for all $t > 0$ if and only if $x = 0$;
- (RN2) $\mu_{\alpha x}(t) = \mu_x(\frac{t}{|\alpha|})$ for all $x \in X$, $\alpha \neq 0$;
- (RN3) $\mu_{x+y}(t+s) \geq T(\mu_x(t), \mu_y(s))$ for all $x, y \in X$ and $t, s \geq 0$.

Every normed space $(X, \|\cdot\|)$ defines a random normed space (X, μ, T_M) , where

$$\mu_x(t) = \frac{t}{t + \|x\|}$$

for all $t > 0$, and T_M is the minimum t -norm. This space is called the *induced random normed space*.

Definition 2.3 ([10]) A *random normed algebra* (X, μ, T, T') is a random normed space (X, μ, T) with algebraic structure such that

- (RN4) $\mu_{xy}(ts) \geq T'(\mu_x(t), \mu_y(s))$ for all $x, y \in X$ and $t, s > 0$, in which T' is a continuous t -norm.

Every normed algebra $(X, \|\cdot\|)$ defines a random normed algebra (X, μ, T_M, T_P) , where

$$\mu_x(t) = \frac{t}{t + \|x\|}$$

for all $t > 0$ if and only if

$$\|xy\| \leq \|x\|\|y\| + s\|y\| + t\|x\|$$

for all $x, y \in X$ and $t, s > 0$. This space is called the *induced random normed algebra*. For more properties and examples of the theory of random normed spaces, we refer to [11–27].

Definition 2.4 Let $(\mathcal{U}, \mu, T, T')$ be a random Banach algebra. Then an *involution* on \mathcal{U} is a mapping $u \rightarrow u^*$ from \mathcal{U} into \mathcal{U} which satisfies the following conditions:

- (1) $u^{**} = u$ for $u \in \mathcal{U}$;
- (2) $(\alpha u + \beta v)^* = \overline{\alpha} u^* + \overline{\beta} v^*$;
- (3) $(uv)^* = v^* u^*$ for $u, v \in \mathcal{U}$.

If, in addition, $\mu_{u^*u}(ts) = T'(\mu_u(t), \mu_u(s))$ for all $u \in \mathcal{U}$ and $t, s > 0$, then \mathcal{U} is a random C^* -algebra.

Following the terminology of [28], a non-empty set G with a ternary operation $[\cdot, \cdot, \cdot] : G \times G \times G \rightarrow G$ is called a *ternary groupoid* and is denoted by $(G, [\cdot, \cdot, \cdot])$. The ternary groupoid $(G, [\cdot, \cdot, \cdot])$ is called *commutative* if $[x_1, x_2, x_3] = [x_{\sigma(1)}, x_{\sigma(2)}, x_{\sigma(3)}]$ for all $x_1, x_2, x_3 \in G$ and all permutations σ of $\{1, 2, 3\}$.

If a binary operation \circ is defined on G such that $[x, y, z] = (x \circ y) \circ z$ for all $x, y, z \in G$, then we say that $[\cdot, \cdot, \cdot]$ is derived from \circ . We say that $(G, [\cdot, \cdot, \cdot])$ is a *ternary semigroup* if the operation $[\cdot, \cdot, \cdot]$ is *associative*, i.e., if

$$[[x, y, z], u, v] = [x, [y, z, u], v] = [x, y, [z, u, v]]$$

for all $x, y, z, u, v \in G$ (see [29]).

A *random C^* -ternary algebra* is a random complex Banach space A , equipped with a ternary product $(x, y, z) \mapsto [x, y, z]$ of A^3 into A , which are \mathbf{C} -linear in the outer variables, conjugate \mathbf{C} -linear in the middle variable, associative in the sense that

$$[x, y, [z, w, v]] = [x, [w, z, y], v] = [[x, y, z], w, v],$$

and satisfies

$$\mu_{[x, y, z]}(tsp) \geq T(\mu_x(t), \mu_y(s), \mu_z(p))$$

and

$$\mu_{[x, x, x]}(t^3) \geq T(\mu_x(t), \mu_x(t), \mu_x(t))$$

(see [28, 30]).

Every random left Hilbert C^* -module is a random C^* -ternary algebra via the ternary product $[x, y, z] := \langle x, y \rangle z$.

If a random C^* -ternary algebra $(A, [\cdot, \cdot, \cdot])$ has the identity, i.e., an element $e \in A$ such that $x = [x, e, e] = [e, e, x]$ for all $x \in A$, then it is routine to verify that A , endowed with $x \circ y := [x, e, y]$ and $x^* := [e, x, e]$, is a unital C^* -algebra. Conversely, if (A, \circ) is a unital C^* -algebra, then $[x, y, z] := x \circ y^* \circ z$ makes A into a C^* -ternary algebra.

A \mathbf{C} -linear mapping $H : A \rightarrow B$ is called a *C^* -ternary algebra homomorphism* if

$$H([x, y, z]) = [H(x), H(y), H(z)]$$

for all $x, y, z \in A$. If, in addition, the mapping H is bijective, then the mapping $H : A \rightarrow B$ is called a *C^* -ternary algebra isomorphism*. A \mathbf{C} -linear mapping $\delta : A \rightarrow A$ is called a C^* -

ternary algebra derivation if

$$\delta([x, y, z]) = [\delta(x), y, z] + [x, \delta(y), z] + [x, y, \delta(z)]$$

for all $x, y, z \in A$ (see [28, 31]).

There are some applications, although still hypothetical, in the fractional quantum Hall effect, the non-standard statistics, supersymmetric theory, and the Yang-Baxter equation (cf. [5, 32, 33]).

Throughout this paper, assume that p, d are non-negative integers with $p + d \geq 3$ and A, B are random C^* -ternary algebras.

Definition 2.5 Let (X, μ, T) be an RN-space.

- (1) A sequence $\{x_n\}$ in X is said to be *convergent* to x in X if, for any $\epsilon > 0$ and $\lambda > 0$, there exists a positive integer N such that $\mu_{x_n-x}(\epsilon) > 1 - \lambda$ whenever $n \geq N$.
- (2) A sequence $\{x_n\}$ in X is called a *Cauchy sequence* if, for any $\epsilon > 0$ and $\lambda > 0$, there exists a positive integer N such that

$$\mu_{x_m-x_n}(\epsilon) > 1 - \lambda$$

whenever $n \geq m \geq N$.

- (3) An RN-space (X, μ, T) is said to be *complete* if every Cauchy sequence in X is convergent to a point in X .

3 Approximation of random C^* -ternary algebras homomorphisms

In this section, we approximate random C^* -ternary algebras homomorphisms of a Cauchy-Jensen additive mapping (see also [34–45]).

For a given mapping $f : A \rightarrow B$, we define

$$\begin{aligned} C_\mu f(x_1, \dots, x_p, y_1, \dots, y_d) \\ := 2f\left(\frac{\sum_{j=1}^p \mu x_j}{2} + \sum_{j=1}^d \mu y_j\right) - \sum_{j=1}^p \mu f(x_j) - 2 \sum_{j=1}^d \mu f(y_j) \end{aligned}$$

for all $\mu \in \mathbf{T}^1 := \{\lambda \in \mathbf{C} : |\lambda| = 1\}$ and $x_1, \dots, x_p, y_1, \dots, y_d \in A$.

One can easily show that a mapping $f : A \rightarrow B$ satisfies

$$C_\mu f(x_1, \dots, x_p, y_1, \dots, y_d) = 0$$

for all $\mu \in \mathbf{T}^1$ and $x_1, \dots, x_p, y_1, \dots, y_d \in A$ if and only if

$$f(\mu x + \lambda y) = \mu f(x) + \lambda f(y)$$

for all $\mu, \lambda \in \mathbf{T}^1$ and $x, y \in A$.

We use the following lemma in this paper.

Lemma 3.1 ([46]) *Let $f : A \rightarrow B$ be an additive mapping such that $f(\mu x) = \mu f(x)$ for all $x \in A$ and $\mu \in \mathbf{T}^1$. Then the mapping f is \mathbf{C} -linear.*

Theorem 3.2 Let r, s , and θ be non-negative real numbers such that $0 < r \neq 1$, $0 < s \neq 3$. Let $\varphi : A^{p+d} \rightarrow D^+$ ($d \geq 2$) and $\psi : A^3 \rightarrow D^+$ such that

$$\varphi_{a(x_1, \dots, x_p, y_1, \dots, y_d)}(t) = \varphi_{x_1, \dots, x_p, y_1, \dots, y_d} \left(\frac{t}{a^r} \right) \quad (1)$$

and

$$\psi_{a(x, y, z)}(t) = \psi_{x, y, z} \left(\frac{t}{a^s} \right) \quad (2)$$

for all $x_1, \dots, x_p, y_1, \dots, y_d, x, y, z \in A$ and $a \in \mathbb{C}$. Suppose that $f : A \rightarrow B$ is a mapping with $f(0) = 0$, satisfying

$$\mu_{C_\mu f(x_1, \dots, x_p, y_1, \dots, y_d)}(t) \geq \varphi_{x_1, \dots, x_p, y_1, \dots, y_d}(t) \quad (3)$$

and

$$\mu_{f([x, y, z]) - [f(x)f(y)f(z)]}(t) \geq \psi_{x, y, z}(t) \quad (4)$$

for all $\mu \in \mathbb{T}^1$, $x_1, \dots, x_p, y_1, \dots, y_d, x, y, z \in A$, and $t > 0$. Then there exists a unique C^* -ternary algebra homomorphism $H : A \rightarrow B$ such that

$$\mu_{f(x) - H(x)}(t) \geq \varphi_{\underbrace{0, \dots, 0}_p, \underbrace{x, \dots, x}_d}(2t(d - d^r)) \quad (5)$$

for all $x \in A$ and $t > 0$.

Proof We prove the theorem when $0 < r < 1$ and $0 < s < 3$. Similarly, one can prove the theorem for other cases. Letting $\mu = 1$, $x_1 = \dots = x_p = 0$, and $y_1 = \dots = y_d = x$ in (3), we get

$$\mu_{2f(dx) - 2df(x)}(t) \geq \varphi_{\underbrace{0, \dots, 0}_p, \underbrace{x, \dots, x}_d}(t) \quad (6)$$

for all $x \in A$ and $t > 0$. If we replace x by $d^n x$ in (6), we get

$$\mu_{\frac{1}{d^{n+1}}f(d^{n+1}x) - \frac{1}{d^n}f(d^n x)}(t) \geq \varphi_{\underbrace{0, \dots, 0}_p, \underbrace{x, \dots, x}_d}(2dtd^{(1-r)n})$$

for all $x \in A$, all non-negative integers n and $t > 0$. Therefore,

$$\mu_{\frac{1}{d^{n+m}}f(d^{n+m}x) - \frac{1}{d^m}f(d^m x)}(t) \geq \varphi_{\underbrace{0, \dots, 0}_p, \underbrace{x, \dots, x}_d} \left(\frac{2dt}{\sum_{k=m}^{m+n} d^{(r-1)k}} \right) \quad (7)$$

for all $x \in A$, non-negative integers n, m and $t > 0$. From this, it follows that the sequence $\{\frac{1}{d^n}f(d^n x)\}$ is a Cauchy sequence for all $x \in A$. Since B is complete, the sequence $\{\frac{1}{d^n}f(d^n x)\}$ converges. Thus one can define the mapping $H : A \rightarrow B$ by

$$H(x) := \lim_{n \rightarrow \infty} \frac{1}{d^n}f(d^n x)$$

for all $x \in A$. Moreover, letting $m = 0$ and passing to the limit $n \rightarrow \infty$ in (7), we get (5). It follows from (3) that

$$\begin{aligned} & \mu_{2H\left(\frac{\sum_{j=1}^p \mu x_j}{2} + \sum_{j=1}^d \mu y_j\right) - \sum_{j=1}^p \mu H(x_j) - 2 \sum_{j=1}^d \mu H(y_j)}(t) \\ &= \lim_{n \rightarrow \infty} \mu_{\frac{1}{d^n} (2f(d^n \frac{\sum_{j=1}^p \mu x_j}{2} + d^n \sum_{j=1}^d \mu y_j) - \sum_{j=1}^p \mu f(d^n x_j) - 2 \sum_{j=1}^d \mu f(d^n y_j))}(t) \\ &\geq \lim_{n \rightarrow \infty} \varphi_{d^n(x_1, \dots, x_p, y_1, \dots, y_d)}(d^n t) \\ &\geq \lim_{n \rightarrow \infty} \varphi_{x_1, \dots, x_p, y_1, \dots, y_d} \left(\frac{d^n}{d^{nr}} t \right) \\ &= 1 \end{aligned}$$

for all $\mu \in \mathbf{T}^1$, $x_1, \dots, x_p, y_1, \dots, y_d \in A$, and $t > 0$. Hence we have

$$2H\left(\frac{\sum_{j=1}^p \mu x_j}{2} + \sum_{j=1}^d \mu y_j\right) = \sum_{j=1}^p \mu H(x_j) + 2 \sum_{j=1}^d \mu H(y_j)$$

for all $\mu \in \mathbf{T}^1$ and $x_1, \dots, x_p, y_1, \dots, y_d \in A$ and so

$$H(\lambda x + \mu y) = \lambda H(x) + \mu H(y)$$

for all $\lambda, \mu \in \mathbf{T}^1$ and $x, y \in A$. Therefore, by Lemma 3.1, the mapping $H : A \rightarrow B$ is \mathbf{C} -linear. It follows from (4) that

$$\begin{aligned} & \mu_{H([x, y, z]) - [H(x), H(y), H(z)]}(t) \\ &= \lim_{n \rightarrow \infty} \mu_{\frac{1}{d^{3n}} (f([d^n x, d^n y, d^n z]) - [f(d^n x), f(d^n y), f(d^n z)])}(t) \\ &= \lim_{n \rightarrow \infty} \mu_{(f([d^n x, d^n y, d^n z]) - [f(d^n x), f(d^n y), f(d^n z)])}(d^{3n} t) \\ &\geq \lim_{n \rightarrow \infty} \psi_{d^n x, d^n y, d^n z}(d^{3n} t) \\ &\geq \lim_{n \rightarrow \infty} \psi_{x, y, z} \left(\frac{d^{3n}}{d^{ns}} \right) = 1 \end{aligned}$$

for all $x, y, z \in A$ and $t > 0$ and so

$$H([x, y, z]) = [H(x), H(y), H(z)]$$

for all $x, y, z \in A$.

Now, let $T : A \rightarrow B$ be another generalized Cauchy-Jensen additive mapping satisfying (5). Then we have

$$\begin{aligned} \mu_{H(x) - T(x)}(t) &= \lim_{n \rightarrow \infty} \mu_{\frac{1}{d^n} (f(d^n x) - T(d^n x))}(t) \\ &= \lim_{n \rightarrow \infty} \mu_{f(d^n x) - T(d^n x)}(d^n t) \\ &\geq \lim_{n \rightarrow \infty} \varphi_{\underbrace{0, \dots, 0}_p, \underbrace{d^n x, \dots, d^n x}_d}(2td^n(d - d^n)) \end{aligned}$$

$$\begin{aligned} &\geq \lim_{n \rightarrow \infty} \varphi_{\underbrace{p}_{0, \dots, 0}, \underbrace{d}_{x_1, \dots, x}} \left(\frac{2td^n(d-d^r)}{d^{nr}} \right) \\ &= 1 \end{aligned}$$

for all $x \in A$ and $t > 0$. So we can conclude that $H(x) = T(x)$ for all $x \in A$. This proves the uniqueness of H . Thus the mapping $H : A \rightarrow B$ is a unique C^* -ternary algebra homomorphism satisfying (5). This completes the proof. \square

Theorem 3.3 *Let $r < 1$, $s < 2$, θ be non-negative real numbers and let $f : A \rightarrow B$ be a mapping satisfying (1), (2), (3) and (4). If there exist a real number $\lambda > 1$ ($0 < \lambda < 1$) and an element $x_0 \in A$ such that*

$$\lim_{n \rightarrow \infty} \frac{1}{\lambda^n} f(\lambda^n x_0) = e' \quad \left(\lim_{n \rightarrow \infty} \lambda^n f\left(\frac{x_0}{\lambda^n}\right) = e' \right),$$

then the mapping $f : A \rightarrow B$ is a C^ -ternary algebra homomorphism.*

Proof By using the proof of Theorem 3.2, there exists a unique C^* -ternary algebra homomorphism $H : A \rightarrow B$ satisfying (5). Now,

$$H(x) = \lim_{n \rightarrow \infty} \frac{1}{\lambda^n} f(\lambda^n x) \quad \left(H(x) = \lim_{n \rightarrow \infty} \lambda^n f\left(\frac{x}{\lambda^n}\right) \right) \quad (8)$$

for all $x \in A$ and all real numbers $\lambda > 1$ ($0 < \lambda < 1$). Therefore, by the assumption, we get that $H(x_0) = e'$. Let $\lambda > 1$ and $\lim_{n \rightarrow \infty} \frac{1}{\lambda^n} f(\lambda^n x_0) = e'$. It follows from (4) and (8) that

$$\begin{aligned} &\mu_{[H(x), H(y), H(z)] - [H(x), H(y), f(z)]}(t) \\ &= \mu_{H[x, y, z] - [H(x), H(y), f(z)]}(t) \\ &= \lim_{n \rightarrow \infty} \mu_{\frac{1}{\lambda^{2n}} (f([\lambda^n x, \lambda^n y, z]) - [f(\lambda^n x), f(\lambda^n y), f(z)])}(t) \\ &= \lim_{n \rightarrow \infty} \mu_{f([\lambda^n x, \lambda^n y, z]) - [f(\lambda^n x), f(\lambda^n y), f(z)]}(\lambda^{2n} t) \\ &\geq \lim_{n \rightarrow \infty} \psi_{\lambda^x, \lambda^y, \lambda^z}(\lambda^{2n} t) \\ &= \psi_{x, y, z}\left(\frac{\lambda^{2n}}{\lambda^{2ns}} t\right) \\ &= 1 \end{aligned}$$

for all $x \in A$ and $t > 0$ and so

$$[H(x), H(y), H(z)] = [H(x), H(y), f(z)]$$

for all $x, y, z \in A$. Letting $x = y = x_0$ in the last equality, we get $f(z) = H(z)$ for all $z \in A$.

Similarly, one can show that $H(x) = f(x)$ for all $x \in A$ when $0 < \lambda < 1$ and $\lim_{n \rightarrow \infty} \lambda^n f\left(\frac{x_0}{\lambda^n}\right) = e'$. Therefore, the mapping $f : A \rightarrow B$ is a C^* -ternary algebra homomorphism. This completes the proof. \square

Theorem 3.4 *Let $r > 1$, $s > 3$, θ be non-negative real numbers and let $f : A \rightarrow B$ be a mapping satisfying (3) and (4). If there exists a real number $0 < \lambda < 1$ ($\lambda > 1$) and an element*

$x_0 \in A$ such that

$$\lim_{n \rightarrow \infty} \frac{1}{\lambda^n} f(\lambda^n x_0) = e' \quad \left(\lim_{n \rightarrow \infty} \lambda^n f\left(\frac{x_0}{\lambda^n}\right) = e' \right),$$

then the mapping $f : A \rightarrow B$ is a C^* -ternary algebra homomorphism.

Proof The proof is similar to the proof of Theorem 3.3 and we omit it. \square

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors carried out the proof. All authors conceived of the study, and participated in its design and coordination. All authors read and approved the final manuscript.

Author details

¹Department of Mathematics Education and the RINS, Gyeongsang National University, Jinju, 660-701, Korea.

²Department of Mathematics, King Abdulaziz University, P.O. Box 80203, Jeddah, Saudi Arabia. ³Department of Mathematics, Iran University of Science and Technology, Tehran, Iran. ⁴Department of Mathematics, Jeju National University, Jeju, 690-756, Korea.

Acknowledgements

The authors would like to thank referees for giving useful comments and suggestions for the improvement of this paper. The third author was supported by the 2015 scientific promotion program funded by Jeju National University.

Received: 22 August 2014 Accepted: 13 December 2014 Published online: 24 January 2015

References

1. Cayley, A: On the 34 concomitants of the ternary cubic. *Am. J. Math.* **4**, 1-15 (1881)
2. Kapranov, M, Gelfand, IM, Zelevinskii, A: *Discriminants, Resultants and Multidimensional Determinants*. Birkhäuser, Berlin (1994)
3. Kerner, R: Ternary algebraic structures and their applications in physics. Univ. P. and M. Curie preprint, Paris (2000)
4. Kerner, R: The cubic chessboard: geometry and physics. *Class. Quantum Gravity* **14**(1A), A203-A225 (1997)
5. Abramov, V, Kerner, R, Le Roy, B: Hypersymmetry: a \mathbb{Z}_3 -graded generalization of supersymmetry. *J. Math. Phys.* **38**, 1650-1669 (1997)
6. Daletskii, YL, Takhtajan, L: Leibniz and Lie algebra structures for Nambu algebras. *Lett. Math. Phys.* **39**, 127-141 (1997)
7. Chang, SS, Cho, YJ, Kang, SM: *Nonlinear Operator Theory in Probabilistic Metric Spaces*. Nova Science Publishers, New York (2001)
8. Schweizer, B, Sklar, A: *Probabilistic Metric Spaces*. North-Holland, New York (1983)
9. Sherstnev, AN: On the notion of a random normed space. *Dokl. Akad. Nauk SSSR* **149**, 280-283 (1963) (in Russian)
10. Mirmostafaei, AK: Perturbation of generalized derivations in fuzzy Menger normed algebras. *Fuzzy Sets Syst.* **195**, 109-117 (2012)
11. Cho, YJ, Rassias, JM, Saadati, R: *Stability of Functional Equations in Random Normed Spaces*. Springer Optimization and Its Applications, vol. 86. Springer, New York (2013)
12. Lee, SJ, Saadati, R: On stability of functional inequalities at random lattice ϕ -normed spaces. *J. Comput. Anal. Appl.* **15**(8), 1403-1412 (2013)
13. Vahidi, J, Park, C, Saadati, R: A functional equation related to inner product spaces in non-Archimedean \mathcal{L} -random normed spaces. *J. Inequal. Appl.* **2012**, 168 (2012)
14. Kang, JI, Saadati, R: Approximation of homomorphisms and derivations on non-Archimedean random Lie C^* -algebras via fixed point method. *J. Inequal. Appl.* **2012**, 251 (2012)
15. Park, C, Eshaghi Gordji, M, Saadati, R: Random homomorphisms and random derivations in random normed algebras via fixed point method. *J. Inequal. Appl.* **2012**, 194 (2012)
16. Rassias, JM, Saadati, R, Sadeghi, G, Vahidi, J: On nonlinear stability in various random normed spaces. *J. Inequal. Appl.* **2011**, 62 (2011)
17. Cho, YJ, Saadati, R: Lattitic non-Archimedean random stability of ACQ functional equation. *Adv. Differ. Equ.* **2011**, 31 (2011)
18. Miheţ, D, Saadati, R: On the stability of the additive Cauchy functional equation in random normed spaces. *Appl. Math. Lett.* **24**(12), 2005-2009 (2011)
19. Miheţ, D, Saadati, R, Vaezpour, SM: The stability of the quartic functional equation in random normed spaces. *Acta Appl. Math.* **110**(2), 797-803 (2010)
20. Mohamadi, M, Cho, YJ, Park, C, Vetro, P, Saadati, R: Random stability on an additive-quadratic-quartic functional equation. *J. Inequal. Appl.* **2010**, Article ID 754210 (2010)
21. Saadati, R, Vaezpour, SM, Cho, YJ: A note to paper 'On the stability of cubic mappings and quartic mappings in random normed spaces'. *J. Inequal. Appl.* **2009**, Article ID 214530 (2009)
22. Chauhan, S, Pant, BD: Fixed point theorems for compatible and subsequentially continuous mappings in Menger spaces. *J. Nonlinear Sci. Appl.* **7**(2), 78-89 (2014)

23. Mihet, D: Common coupled fixed point theorems for contractive mappings in fuzzy metric spaces. *J. Nonlinear Sci. Appl.* **6**(1), 35-40 (2013)
24. Park, C, Eshaghi Gordji, M, Najati, A: Generalized Hyers-Ulam stability of an AQCC-functional equation in non-Archimedean Banach spaces. *J. Nonlinear Sci. Appl.* **3**(4), 272-281 (2010)
25. Shakeri, S: A contraction theorem in Menger probabilistic metric spaces. *J. Nonlinear Sci. Appl.* **1**(3), 189-193 (2008)
26. Shakeri, S: A note on the 'A contraction theorem in Menger probabilistic metric spaces'. *J. Nonlinear Sci. Appl.* **2**(1), 25-26 (2009)
27. Zaharia, C: On the probabilistic stability of the monomial functional equation. *J. Nonlinear Sci. Appl.* **6**(1), 51-59 (2013)
28. Amyari, M, Moslehian, MS: Approximately ternary semigroup homomorphisms. *Lett. Math. Phys.* **77**, 1-9 (2006)
29. Bazunova, N, Borowiec, A, Kerner, R: Universal differential calculus on ternary algebras. *Lett. Math. Phys.* **67**, 195-206 (2004)
30. Zettl, H: A characterization of ternary rings of operators. *Adv. Math.* **48**, 117-143 (1983)
31. Moslehian, MS: Almost derivations on C^* -ternary rings. *Bull. Belg. Math. Soc. Simon Stevin* **14**, 135-142 (2007)
32. Kerner, R: Ternary algebraic structures and their applications in physics. Preprint
33. Vainerman, L, Kerner, R: On special classes of n -algebras. *J. Math. Phys.* **37**, 2553-2565 (1996)
34. Ulam, SM: A Collection of the Mathematical Problems. Interscience, New York (1960)
35. Hyers, DH: On the stability of the linear functional equation. *Proc. Natl. Acad. Sci. USA* **27**, 222-224 (1941)
36. Aoki, T: On the stability of the linear transformation in Banach spaces. *J. Math. Soc. Jpn.* **2**, 64-66 (1950)
37. Rassias, TM: On the stability of the linear mapping in Banach spaces. *Proc. Am. Math. Soc.* **72**, 297-300 (1978)
38. Rassias, TM: Problem 16; 2. Report of the 27th International Symp. on Functional Equations. *Aequ. Math.* **39**, 292-293 (1990)
39. Gajda, Z: On stability of additive mappings. *Int. J. Math. Math. Sci.* **14**, 431-434 (1991)
40. Rassias, TM, Šemrl, P: On the behaviour of mappings which do not satisfy Hyers-Ulam stability. *Proc. Am. Math. Soc.* **114**, 989-993 (1992)
41. Găvruta, P: A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings. *J. Math. Anal. Appl.* **184**, 431-436 (1994)
42. Jung, S: On the Hyers-Ulam-Rassias stability of approximately additive mappings. *J. Math. Anal. Appl.* **204**, 221-226 (1996)
43. Czerwik, P: Functional Equations and Inequalities in Several Variables. World Scientific, Singapore (2002)
44. Hyers, DH, Isac, G, Rassias, TM: Stability of Functional Equations in Several Variables. Birkhäuser, Basel (1998)
45. Isac, G, Rassias, TM: Stability of ψ -additive mappings: applications to nonlinear analysis. *Int. J. Math. Math. Sci.* **19**, 219-228 (1996)
46. Park, C: Homomorphisms between Poisson JC^* -algebras. *Bull. Braz. Math. Soc.* **36**, 79-97 (2005)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com